
SCIENTIFIC AMERICAN™

BIOLOGY

Devastating Wheat Fungus Appears in Asia for the First Time

Scientists race to determine the origin of a Bangladesh outbreak, which they warn could spread farther afield

By Ewen Callaway, Nature magazine on April 27, 2016



Credit: tgraham/Flickr, CC BY-SA 2.0

Fields are ablaze in Bangladesh, as farmers struggle to contain Asia's first outbreak of a fungal disease that periodically devastates crops in South

America. Plant pathologists warn that wheat blast could spread to other parts of south and southeast Asia, and are hurrying to trace its origins.

“It’s important to know what the strain is,” says Sophien Kamoun, a biologist at the Sainsbury Laboratory in Norwich, UK, who has created a website, Open Wheat Blast (go.nature.com/bkczwf), to encourage researchers to share data.

Efforts are also under way to find wheat genes that confer resistance to the disease.

First detected in February and confirmed with genome sequencing by Kamoun’s lab this month, the wheat-blast outbreak has already caused the loss of more than 15,000 hectares of crops in Bangladesh. “It’s really an explosive, devastating disease,” says plant pathologist Barbara Valent of Kansas State University in Manhattan, Kansas. “It’s really critical that it be controlled in Bangladesh.”

After rice, wheat is the second most cultivated grain in Bangladesh, which has a population of 156 million people. More broadly, inhabitants of south Asia grow 135 million tonnes of wheat each year.

Wheat blast is caused by the fungus *Magnaporthe oryzae*. Since 1985, when scientists discovered it in Brazil’s Paraná state, the disease has raced across South America.

The fungus is better known as a pathogen of rice. But unlike in rice, where *M. oryzae* attacks the leaves, the fungus strikes the heads of wheat, which are difficult for fungicides to reach. A 2009 outbreak in wheat cost Brazil one-third of that year’s crop. “There are regions in South America where they don’t grow wheat because of the disease,” Valent says. Wheat blast was spotted in Kentucky in 2011, but vigorous surveillance helped to stop it spreading in the

United States.

In South America, the disease tends to take hold in hot and humid spells. Such conditions are present in Bangladesh, and the disease could migrate across south and southeast Asia, say plant pathologists. In particular, it could spread over the Indo-Gangetic Plain through Bangladesh, northern India and eastern Pakistan, warn scientists at the Bangladesh Agricultural Research Institute (BARI) in Nashipur.

Bangladeshi officials are burning government-owned wheat fields to contain the fungus, and telling farmers not to sow seeds from infected plots. The BARI hopes to identify wheat varieties that are more tolerant of the fungus and agricultural practices that can keep it at bay, such as crop rotation and seed treatment.

It is unknown how wheat blast got to Bangladesh. One possibility is that a wheat-infecting strain was brought in from South America, says Nick Talbot, a plant pathologist at the University of Exeter, UK. Another is that an *M. oryzae* strain that infects south Asian grasses somehow jumped to wheat, perhaps triggered by an environmental shift: that is what happened in Kentucky, when a rye-grass strain infected wheat.

To tackle the question, this month Kamoun's lab sequenced a fungus sample from Bangladesh. The strain seems to be related to those that infect wheat in South America, says Kamoun, but data from other wheat-infecting strains and strains that plague other grasses are needed to pinpoint the outbreak's origins conclusively.

The Open Wheat Blast website might help. Kamoun has uploaded the Bangladeshi data, and Talbot has deposited *M. oryzae* sequences from wheat in Brazil. Talbot hopes that widely accessible genome data could help to combat

the outbreak. Researchers could use them to screen seeds for infection or identify wild grasses that can transmit the fungus to wheat fields.

Rapid data sharing is becoming more common in health emergencies, such as the outbreak of Zika virus in the Americas. Kamoun and Talbot say that their field should follow suit. “The plant-pathology community has a responsibility to allow data to be used to combat diseases that are happening now, and not worry too much about whether they may or may not get a *Nature* paper out of it,” says Talbot.

Last month, Valent’s team reported the first gene variant known to confer wheat-blast resistance (C. D. Cruz *et al. Crop Sci.* <http://doi.org/bfk7>; 2016), and field trials of crops that bear the resistance gene variant have begun in South America. But plant pathologists say that finding one variant is not enough: wheat strains must be bred with multiple genes for resistance, to stop *M. oryzae* quickly overcoming their defences.

The work could help in the Asian crisis, says Talbot. “What I would hope for out of this sorry situation,” he says, “is that there will be a bigger international effort to identify resistance genes.”

This article is reproduced with permission and was first published on April 27, 2016.



ADVERTISEMENT

ABOUT THE AUTHOR(S)

Scientific American is part of Springer Nature, which owns or has commercial relations with thousands of scientific publications (many of them can be found at www.springernature.com/us). Scientific American maintains a strict policy of editorial independence in reporting developments in science to our readers.

© 2016 SCIENTIFIC AMERICAN, A DIVISION OF NATURE AMERICA, INC.

ALL RIGHTS RESERVED.