

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/259385648>

Emergence of a novel drug resistant H7N9 influenza virus: Evidence based clinical potential of a natural IFN- α for infection control and treatment

ARTICLE *in* EXPERT REVIEW OF ANTI-INFECTIVE THERAPY · DECEMBER 2013

Impact Factor: 3.46 · DOI: 10.1586/14787210.2014.870885 · Source: PubMed

CITATIONS

5

READS

95

7 AUTHORS, INCLUDING:



[Qinfang Liu](#)

Kansas State University

34 PUBLICATIONS 356 CITATIONS

[SEE PROFILE](#)



[William Marvin Mitchell](#)

Vanderbilt University

178 PUBLICATIONS 5,031 CITATIONS

[SEE PROFILE](#)



[Wenjun Ma](#)

Kansas State University

56 PUBLICATIONS 1,476 CITATIONS

[SEE PROFILE](#)



[Jürgen A Richt](#)

Kansas State University

12 PUBLICATIONS 117 CITATIONS

[SEE PROFILE](#)

Emergence of a novel drug resistant H7N9 influenza virus: evidence based clinical potential of a natural IFN- α for infection control and treatment

Expert Rev. Anti Infect. Ther. 12(2), 000–000 (2014)

Qinfang Liu¹,
Jingjiao Ma¹,
David R Strayer²,
William M Mitchell³,
William A Carter²,
Wenjun Ma¹ and
Juergen A Richt*¹

¹Center of Excellence for Emerging and Zoonotic Animal Diseases (CEEZAD), Kansas State University, Manhattan, KS 66506-5601, USA

²Hemispherx Biopharma, Inc., Philadelphia, PA, USA

³Vanderbilt University, Nashville, TN, USA

*Author for correspondence:
Tel.: +1 785 532 2793
Fax: +1 785 532 3373
jricht@ksu.edu

The novel avian H7N9 influenza virus has caused more than 130 human infections with 43 deaths (as of September, 2013) in China. Because of the lack of existing immunity against H7 subtype influenza viruses in the human population and the absence of a licensed commercial vaccine, antiviral drugs are critical tools for the treatment of infection with this novel H7N9. Both M2-ion channel blockers and neuraminidase inhibitors are used as antiviral drugs for influenza infections of humans. The emerging H7N9 viruses are resistant to the M2-ion channel blockers because of a S31N mutation in the M2 protein; additionally, some H7N9 isolates have gained neuraminidase R292K substitution resulting in broad resistance to neuraminidase inhibitors. In this study we report that Alferon N can inhibit wild type and 292K H7N9 viruses replication *in vitro*. Since Alferon N is approved for clinical use, this would allow a rapid regulatory approval process for this drug under pandemic threat.

KEYWORDS: Alferon N • antiviral drug • H7N9 • influenza • neuraminidase inhibitors • oseltamivir resistance • tamiflu

Emergence of the novel H7N9

A new deadly influenza A H7N9 strain of avian influenza virus, first identified in China in March 2013 with human and avian cases, has been observed in various eastern and inland Chinese provinces [1,2]. There have been a total of 132 confirmed cases with 43 deaths with people over 60 appearing to be more susceptible to the H7N9-associated severe respiratory disease [3–6]. Patients infected by H7N9 have exhibited viral pneumonia, acute kidney failure, acute respiratory distress syndrome, diffuse intravascular coagulation and septic shock. Although the H7N9 subtype can be highly virulent in humans, it exhibits low pathogenicity in avian hosts, perhaps due in part to the absence of a multibasic amino acid sequence at the hemagglutinin cleavage site [1,7,8]. Infected chickens, however, can shed virus very

efficiently via the oropharynx, and we and others have observed limited shedding by the cloacal route despite no clinical signs of infection [9]. This unusual low pathogenicity of the avian H7N9 viruses makes an H7N9 infection in avian species hard to detect and monitor. Experts remain fearful of the potential for this virus to mutate into a form easily transmissible between humans, although to date, there has been no clear confirmation of human-to-human transmission [4].

Genetic analysis shows that the novel H7N9 virus is a triple reassortant virus carrying genes from H7N3, H9N2 and H11N9 or H2N9 avian influenza A viruses. It is assumed that the novel H7N9 subtype virus was generated by a single or double independent reassortment event in ducks or chickens in China [10]. Most of the H7N9 viruses carry certain mammalian adaptation signatures (i.e.,

Table 1. Genetic signature on various gene products of Anhui/1/H7N9 and Shanghai/1/H7N9 viruses.

Virus	PB2		HA		NA	M2	
	627	701	226 [†]	228 [†]	292 [‡]	27	31
Anhui/1/H7N9	K	D	L	G	R [§]	V	N
Shanghai/1/H7N9	K	D	Q	G	K [¶]	V	N

[†]Positions 226, 228 are H3 numbering system.
[‡]Position 292 is N2 numbering system.
[§]NA292R affords oseltamivir sensitivity.
[¶]NA292K affords oseltamivir resistance.

PB2-627K, HA-226L), which have been proved critical in mammalian adaptation and transmission based on previous studies, [11], and especially the 226L in the H7 HA protein. Virulence and transmissibility of this novel H7N9 influenza virus have been characterized in different mammalian animal models, including mice, ferrets, macaques and pigs [7,9,12–14]. In ferrets, the H7N9 virus showed transmission by direct contact among ferrets and limited transmissibility by respiratory droplets [7,12–14]. Moreover, the novel H7N9 virus replicated efficiently in both upper and lower respiratory tracts of infected nonhuman primates (*Cynomolgus macaques*). Emergent H7N9 viruses isolated from recent human cases have increased infectivity and lethality in mice compared with avian H9N2 viruses and genetically related H7N9 viruses isolated from wild birds before 2013 [7,12]. Swine are an important intermediate host for influenza viruses [15]; one of the concerns is that H7N9 viruses may jump into pigs, they might replicate efficiently in this host, adapt to the mammalian receptor repertoire and reassort with endemic swine influenza viruses. Previous studies showed that pigs infected intranasally with H7N9 viruses did not transmit virus to contact pigs [14], while pigs infected intratracheally with H7N9 can transmit to sentinel animals with low efficiency (Liu *et al.*: submitted). These facts warrant increased surveillance for H7N9 influenza viruses in pigs.

Antivirals & Tamiflu[®] resistance of the novel H7N9 virus

The novel H7N9 virus rapidly spread over a large geographical region in China after the first reported case in March 2013 with the live poultry markets thought to be the source of human infections, suggesting that H7N9 viruses might be silently prevalent in poultry in China [10]. Continuing detection in poultry and sporadic human infections raises concerns on the emergence of a H7N9 virus, which can efficiently transmit between humans. Vaccination is a very efficient way to prevent influenza infection; however, no commercial vaccine specific for H7N9 is currently available. Moreover, humans lack any immunological memory against H7N9, which is another concern of its potential to result in a pandemic if easy human-to-human transmission is acquired.

Antiviral treatment is an important strategy in helping to control and prevent influenza infections. As recommended by the CDC in Atlanta, USA, antiviral treatment against influenza

infections should be applied as early as possible, ideally within 48 h after onset of illness. Antiviral drugs are readily available to treat influenza infections including this novel H7N9 virus. For influenza infections, M2-ion channel blockers (e.g., amantadine and rimantadine) and neuraminidase inhibitors (e.g., oseltamivir, zanamivir and peramivir) have been mainly used as antiviral drugs, and these compounds are approved by the US FDA [16,17]. However, the H7N9 is resistant to both amantadine and rimantadine because an S31N mutation exists in the M2 protein, which confers M2-ion channel blocker resistance [18]. Therefore, the neuraminidase inhibitors are the only FDA-approved compounds that can be used for the treatment of the novel H7N9 infection and other M2-ion channel blocker-resistant influenza viruses [19].

Clinical studies show that oseltamivir treatment reduces the viral load in the respiratory tract of patients with H7N9 infection [5]. However, some patients with oseltamivir treatment failure had a poor clinical outcome. Further studies on the H7N9 viruses isolated from those patients showed that an R292K substitution occurred in the NA protein; this NA change is associated with resistance to all neuraminidase inhibitors including oseltamivir (Tamiflu[®]) [20]. Surprisingly, in humans treated with oseltamivir, the R292K substitution occurred as early as 2 days post treatment with the 292K genotype becoming the predominant virus population at 7 days post treatment. Neuraminidase inhibition assays using the resistant H7N9 virus showed 100- and 30-fold reduced susceptibility to the two FDA-approved inhibitors, oseltamivir and zanamivir, respectively [5]. Subsequently, neuraminidase inhibitor resistance of H7N9 viruses in humans resulted in antiviral treatment failure and a poor clinical outcome. Further studies confirmed that H7N9 isolates with a dominant 292K genotype are resistant to zanamivir, peramivir and oseltamivir although this resistance can be masked by a mixed R/K genotype on position 292 of the NA protein in a viral quasispecies population [21]. Rapid emergence of antiviral drug-resistant mutants after treatment with oseltamivir or zanamivir is raising concerns about the prevalence of such resistant H7N9 viruses in the humans and other mammalian populations. This emphasizes the importance of the development of new antiviral influenza drugs or the use of already developed FDA-approved drugs for other infectious diseases, which might also work against influenza. Other influenza antivirals are under development, including thiazides that are anti-influenza molecules targeting the viral hemagglutinin at the post-translational level [22]; Favipiravir (T-705), a novel viral RNA polymerase inhibitor [23] and Aprotinin, a protease inhibiting the cleavage of the HA protein [24]. None of these compounds, however, have been approved for clinical treatment for influenza infection.

Alferon N inhibits replication of oseltamivir resistant & sensitive H7N9 viruses in vitro

Type I interferons (IFN), including IFN- α and IFN- β produced by host cells in response to the presence of viral pathogens, play a critical role in pathogen clearance during infection [25]. Previous studies showed that the 2009 pandemic




Figure 1. Sensitivity of human H7N9 influenza viruses to Tamiflu (A & B) and Alferon N (C & D) *in vitro*. A549 cells were infected with Shanghai/1/H7N9 (containing K at NA residue 292) or Anhui/1/H7N9 (containing R at NA residue 292) viruses at MOI of 0.01. Supernatants were collected at 12, 24, 36 and 48 h post infection and were titrated in MDCK cells, and all studies were conducted in BSL3/BSL3Ag laboratory located at Biosecurity Research Institute (BSL3/BSL3Ag) located at Kansas State University. All data were analyzed by using analysis of variance in GraphPad Prism version 5.0 (GraphPad software Inc, CA, USA); $p < 0.05$ was considered statistically significant. Bars represent the mean values \pm SEM for three independent replicates. * $p < 0.05$; ** $p < 0.01$.

H1N1 and the highly pathogenic H5N1 viruses are highly sensitive to the antiviral actions of type I IFNs [26,27]. Type I IFN upregulates the expression of several IFN-stimulated antiviral gene products such as protein kinase R (PKR), 2',5'-oligoadenylate synthetase and the Mx protein [28]. The activated protein kinase R phosphorylates the eukaryotic initiation factor eIF2 α , thereby oligoadenylate synthetase inhibits viral protein synthesis [28]; and the Mx protein interacts with viral components to block generation of new virus particles [26,30].

Alferon N (IFN- α -n3) is a natural IFN- α product derived from human leukocytes. In previous studies using severe acute respiratory syndrome (SARS) virus, Alferon N was the only IFN-based drug to be active against SARS virus at clinically achievable serum levels with the failure of other tested IFN-based clinical products [31]. Additionally, Alferon N has a better safety profile than recombinant IFNs [32]. We tested antiviral activity of Alferon N against the novel H7N9 virus in human alveolar epithelial cells (A549) using two human H7N9 isolates: A/Anhui/1/2013 (Anhui/1/H7N9) and

A/Shanghai/1/2013 (Shanghai/1/H7N9). Both viruses were isolated from human cases in China. Sequence information showed that the Anhui/1/H7N9 contains the NA-292R residue and is sensitive to NA inhibitors; in contrast, the Shanghai/1/H7N9 contains a NA-292K residue that is associated with resistance to NA inhibitors. The genetic signatures of both viruses are summarized in TABLE 1. Oseltamivir (Tamiflu®), an NA inhibitor, was tested on A549 cell as a well-known and widely used anti-influenza drug.

Confluent A549 cells in 48-well plates were treated with different units of Alferon N (10,000, 1,000 and 100 IU/ml obtained from Hemispherx Biopharma) for 4 h before infection and untreated cells served as controls; thereafter, both Alferon N treated and untreated A549 cells were infected with the Anhui/1/H7N9 and Shanghai/1/H7N9 viruses at a multiplicity of infection of 0.01 under BSL3/BSL3Ag conditions. Combination of a short natural IFN- α preincubation and a multiplicity of infection of 0.01 simulate the conditions of an *in vivo* natural infection. A relatively small number of cells are

180 initially infected resulting in the induction of type 1 IFNs that
in turn induce innate immune responses in uninfected cells. The
supernatants were collected at 12, 24, 36 and 48 h post infection
and titrated for virus content on Madin-Darby canine kidney
(MDCK) cells in 96-well plates as described previously [33]. For
185 comparison, Oseltamivir (Tamiflu[®] obtained from Roche) was
tested against both H7N9 viruses on A549 cells using a concen-
tration of 0.3, 0.2, 0.1 and 0.01 microgram/ml with treatment
starting at the time of infection.

The results showed that oseltamivir (Tamiflu[®]) was able to
190 inhibit the replication of NA inhibitor sensitive Anhui/1/
H7N9 virus at 36 and 48 h post infection at a concentration
ranging from 0.1 to 0.3 microgram/ml (FIGURE 1A); in contrast,
oseltamivir (Tamiflu[®]) treatment with the same doses did not
inhibit the NA inhibitor-resistant Shanghai/1/H7N9 virus
195 (FIGURE 1B), confirming that the 292K residue in NA of Shang-
hai/1/H7N9 conferred resistance to oseltamivir. Antiviral activi-
ty of Alferon N was found against both the Shanghai/1/
H7N9 and Anhui/1/H7N9; clear inhibition of virus replication
was found at 36 and 48 h post infection (FIGURE 1C & D). Alferon
200 N with a concentration of 10,000 IU/ml significantly reduced
Anhui/1/H7N9 virus load at 36 h post infection, and all three
doses (100, 1,000 and 10,000 IU/ml) significantly inhibit repli-
cation of oseltamivir sensitive Anhui/1/H7N9 and oseltamivir-
resistant Shanghai/1/H7N9 at 48 h post infection (FIGURE 1C &
205 D). Our data demonstrate that Alferon N is effective against
both oseltamivir sensitive and resistant H7N9 viruses and pro-
vide sufficient evidence to justify the costly *in vivo* studies
(BSL3/BSL3Ag) required for regulatory approval.

Expert commentary & five-year view

210 The emergence of NA inhibitor-resistant strains of the novel
H7N9 virus in treated patients requires alternative antiviral
compounds for human therapy, which will be critical in case of
a potential pandemic. Our results show that both oseltamivir
(Tamiflu[®]) and Alferon N have significant inhibitory effect on
215 the NA inhibitor sensitive H7N9 virus. In contrast, Alferon N,
not Tamiflu, had an inhibitory effect on the neuraminidase-
resistant Shanghai/1/H7N9 virus. These results provide evi-
dence for a new therapeutic strategy to mitigate the public
health hazards associated with H7N9 infection in humans and
220 the potential spread of NA inhibitor-resistant H7N9 and other
influenza viruses of potential high virulence for humans.

The emergence of an influenza pandemic is unpredictable.
No universal influenza vaccine is available so far. The licensed
anti-influenza drugs (M2-ion channel blockers and neurami-
nidase inhibitors) are limited in clinical treatment of infected
225 patients because of rapid emergence of drug-resistant muta-
tions in the targeted genes. One of the concerns of a poten-
tial influenza pandemic is high mortality (e.g., H5N1,
H7N9). The rapid disease progression characteristic of pan-
demic influenza viruses and the emergence of resistance
230 to inhibitors of viral function will require other approaches to
limit or terminate disease and spread the virus. In addition to
the rapid regulatory approval of Alferon N or other anti-
influenza drugs for this indication, the development of novel
anti-influenza drugs using different mechanisms or targeting
235 different genes (e.g., polymerase genes, nucleoprotein gene
and hemagglutinin gene, etc.) innate immune responses, and/
or combination therapy is necessary for better pandemic
defense.

Acknowledgements

240 We would like to thank RJ Webby from St Jude Children's Research Hos-
pital for supplying us with the A/Anhui/1/2013 virus stock and A García-
Sastre from Icahn School of Medicine at Mount Sinai for supplying A/
Shanghai/1/2013 virus stock. We would also like to thank Haixia Liu
and laboratory coordinators from Biosecurity Research Institute (BSL3/
245 BSL3Ag) located at Kansas State for providing technical support and NM
Bouvier for cell culture compatible Tamiflu[®].

Financial & competing interests disclosure

250 This work was partially funded by the National Institute of Allergy and
Infectious Diseases, National Institutes of Health, Department of Health
and Human Services, under contract numbers HHSN266200700005C,
by the U.S. Department of Homeland Security under Grant Award Num-
ber 2010-ST-AG0001, and by Hemispherx Biopharma Inc (HEB).
Q Liu is a scientific adviser to Hemispherx Biopharma (HEB); JA Richt
255 is a scientific adviser and received funds from HEB; DR Strayer and WA
Carter are management employees and shareholders of HEB; WM Mitch-
ell is an independent member of HEB BOD and HEB shareholder. The
authors have no other relevant affiliations or financial involvement with
any organization or entity with a financial interest in or financial conflict
260 with the subject matter or materials discussed in the manuscript apart
from those disclosed.

No writing assistance was utilized in the production of this manuscript.

Key issues

- Antiviral treatment is an important strategy in controlling and preventing influenza viral infections. Current pharmaceuticals approved for the prophylaxis and/or treatment of influenza inhibit the function of the neuraminidase and M2-ion channel viral proteins.
- High mutational rates are characteristic of influenza viruses, resulting in viruses with resistance to neuraminidase or M2 inhibitors. Such viruses present a high-risk human public health threat for emerging avian influenza viruses with high pathogenicity for humans such as H7N9.
- Some H7N9 isolates have gained resistance to neuraminidase inhibitors. Alferon N (natural interferon) has broad anti-viral activities and inhibits the replication of both wild-type and neuraminidase inhibitor-resistant H7N9 viruses.
- An imminent pandemic threat from H7N9 or other avian influenza viruses with resistance to neuraminidase and M2 inhibitor demands research on novel anti-influenza drugs targeting different viral genes and/or innate immune responses.

References

Papers of special note have been highlighted as:

• of interest

•• of considerable interest

- Gao R, Cao B, Hu Y, et al. Human infection with a novel avian-origin influenza A (H7N9) virus. *N Engl J Med* 2013; 368(20):1888-97
- Detailed introduction to human infection with novel H7N9 in China.**
- Mei Z, Lu S, Wu X, et al. Avian influenza A(H7N9) virus infections, Shanghai, China. *Emerg Infect Dis* 2013;19(7):1179-81
- Salathe M, Freifeld CC, Mekaru SR, et al. Influenza A (H7N9) and the importance of digital epidemiology. *N Engl J Med* 2013; 369(5):401-4
- Li Q, Zhou L, Zhou M, et al. Preliminary report: epidemiology of the Avian Influenza A (H7N9) outbreak in China. *N Engl J Med* 2013; Epub ahead of print
- Hu Y, Lu S, Song Z, et al. Association between adverse clinical outcome in human disease caused by novel influenza A H7N9 virus and sustained viral shedding and emergence of antiviral resistance. *Lancet* 2013;381(9885):2273-9
- Detailed description of drug-resistant H7N9 infection in patients.**
- To KK, Chan JF, Chen H, et al. The emergence of influenza A H7N9 in human beings 16 years after influenza A H5N1: a tale of two cities. *Lancet Infect Dis* 2013; 13(9):809-21
- Review of H7N9 and H5N1 viruses.**
- Zhang Q, Shi J, Deng G, et al. H7N9 influenza viruses are transmissible in ferrets by respiratory droplet. *Science* 2013; 341(6144):410-14
- Zhou J, Wang D, Gao R, et al. Biological features of novel avian influenza A (H7N9) virus. *Nature* 2013;499(7459):500-3
- Watanabe T, Kiso M, Fukuyama S, et al. Characterization of H7N9 influenza A viruses isolated from humans. *Nature* 2013;501(7468):551-5
- Lam TT, Wang J, Shen Y, et al. The genesis and source of the H7N9 influenza viruses causing human infections in China. *Nature* 2013;502(7470):241-4
- Hatta M, Gao P, Halfmann P, Kawaoka Y. Molecular basis for high virulence of Hong Kong H5N1 influenza A viruses. *Science* 2001;293(5536):1840-2
- Belser JA, Gustin KM, Pearce MB, et al. Pathogenesis and transmission of avian influenza A (H7N9) virus in ferrets and mice. *Nature* 2013;501(7468):556-9
- Richard M, Schrauwen EJ, de Graaf M, et al. Limited airborne transmission of H7N9 influenza A virus between ferrets. *Nature* 2013;501(7468):560-3
- Zhu H, Wang D, Kelvin DJ, et al. Infectivity, transmission, and pathology of human-isolated H7N9 influenza virus in ferrets and pigs. *Science* 2013;341(6142): 183-6
- Ma W, Kahn RE, Richt JA. The pig as a mixing vessel for influenza viruses: human and veterinary implications. *J Mol Genet Med* 2008;3(1):158-66
- Schmidt AC. Antiviral therapy for influenza: a clinical and economic comparative review. *Drugs* 2004;64(18):2031-46
- Beigel J, Bray M. Current and future antiviral therapy of severe seasonal and avian influenza. *Antiviral Res* 2008;78(1):91-102
- Grambas S, Bennett MS, Hay AJ. Influence of amantadine resistance mutations on the pH regulatory function of the M2 protein of influenza A viruses. *Virology* 1992; 191(2):541-9
- Hay AJ, Hayden FG. Oseltamivir resistance during treatment of H7N9 infection. *Lancet* 2013;381(9885):2230-2
- Yen HL, Herlocher LM, Hoffmann E, et al. Neuraminidase inhibitor-resistant influenza viruses may differ substantially in fitness and transmissibility. *Antimicrob Agents Chemother* 2005;49(10):4075-84
- Yen HL, McKimm-Breschkin JL, Choy KT, et al. Resistance to neuraminidase inhibitors conferred by an R292K mutation in a human influenza virus H7N9 isolate can be masked by a mixed R/K viral population. *MBio* 2013;4(4):e00396-13
- Rosignol JF, La Frazia S, Chiappa L, et al. Thiazolidines, a new class of anti-influenza molecules targeting viral hemagglutinin at the post-translational level. *J Biol Chem* 2009;284(43):29798-808
- Furuta Y, Gowen BB, Takahashi K, et al. Favipiravir (T-705), a novel viral RNA polymerase inhibitor. *Antiviral Res* 2013;100(2):446-54
- Zhirnov OP, Klenk HD, Wright PF. Aprotinin and similar protease inhibitors as drugs against influenza. *Antiviral Res* 2011; 92(1):27-36
- Antonelli G. Biological basis for a proper clinical application of alpha interferons. *New Microbiol* 2008;31(3):305-18
- Review of biological basis of alpha interferons.**
- Haasbach E, Droebner K, Vogel AB, Planz O. Low-dose interferon Type I treatment is effective against H5N1 and swine-origin H1N1 influenza A viruses in vitro and in vivo. *J Interferon Cytokine Res* 2011;31(6):515-25
- Osterlund P, Pirhonen J, Ikonen N, et al. Pandemic H1N1 2009 influenza A virus induces weak cytokine responses in human macrophages and dendritic cells and is highly sensitive to the antiviral actions of interferons. *J Virol* 2010;84(3):1414-22
- Silverman RH. Viral encounters with 2',5'-oligoadenylate synthetase and RNase L during the interferon antiviral response. *J Virol* 2007;81(23):12720-9
- Garcia MA, Gil J, Ventoso I, et al. Impact of protein kinase PKR in cell biology: from antiviral to antiproliferative action. *Microbiol Mol Biol Rev* 2006;70(4): 1032-60
- Haller O, Staeheli P, Kochs G. Interferon-induced Mx proteins in antiviral host defense. *Biochimie* 2007;89(6-7):812-18
- Tan EL, Ooi EE, Lin CY, et al. Inhibition of SARS coronavirus infection in vitro with clinically approved antiviral drugs. *Emerg Infect Dis* 2004;10(4):581-6
- Strayer DR, Carter WA. Recombinant and natural human interferons: analysis of the incidence and clinical impact of neutralizing antibodies. *J Interferon Cytokine Res* 2004; 32(3):95-102
- Review of clinical outcomes following treatment of a wide range of diseases with recombinant interferons and natural interferons.**
- Liu Q, Qiao C, Marjuki H, et al. Combination of PB2 271A and SR polymorphism at positions 590/591 is critical for viral replication and virulence of swine influenza virus in cultured cells and in vivo. *J Virol* 2012;86(2):1233-7

Notice of correction

Information on page 4 of the article has been changed post publication and 'mg/ml' had been changed to 'microgram/ml'.